

**Wilcoxon's test: Versions in {3,...,9} (R<sup>+</sup>) vs. versions in {2,...,7} (R<sup>-</sup>) on Tst. and R**

| Comparison                  | Measure | R <sup>+</sup> | R <sup>-</sup> | Hypothesis ( $\alpha = 0.05$ ) | p-value |
|-----------------------------|---------|----------------|----------------|--------------------------------|---------|
| GR-MF(3-9) vs. GR-MF(2-7)   | Tst.    | 14             | <b>64</b>      | Rejected                       | 0.049   |
| GA-WM(3-9) vs. GA-WM(2-7)   | Tst.    | 13             | <b>65</b>      | Rejected                       | 0.041   |
| GLD-WM(3-9) vs. GLD-WM(2-7) | Tst.    | 32             | <b>46</b>      | Accepted                       | 0.583   |
| GR-MF(3-9) vs. GR-MF(2-7)   | R       | 10             | <b>68</b>      | Rejected                       | 0.023   |
| GA-WM(3-9) vs. GA-WM(2-7)   | R       | 0              | <b>78</b>      | Rejected                       | 0.002   |
| GLD-WM(3-9) vs. GLD-WM(2-7) | R       | 0              | <b>78</b>      | Rejected                       | 0.002   |